

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

(NASA-CP-176242) COMETARY ATMOSPHERES:
MODELING THE SPATIAL DISTRIBUTION OF
OBSERVED NEUTRAL RADICALS Interim Report,
Jun. - Aug. 1985 (Atmospheric and
Environmental Research) 8 p HC A02/MF A01

N86-11133

Unclas
27532

G3/90

Cometary Atmospheres:
Modeling the Spatial Distribution
of Observed Neutral Radicals

Michael R. Combi

Atmospheric and Environmental Research, Inc.
840 Memorial Drive
Cambridge, MA 02139

October 1985

Interim Report for the Period
June 1, 1985 to August 31, 1985

Prepared for
NASA Headquarters



TECHNICAL REPORT STANDARD TITLE PAGE

1. Report No.	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle Cometary Atmospheres: Modeling the Spatial Distribution of Observed Neutral Radicals		5. Report Date October 1985	
		6. Performing Organization Code	
7. Author(s) Michael R. Combi		8. Performing Organization Report No.	
9. Performing Organization Name and Address Atmospheric and Environmental Research, Inc. 840 Memorial Drive Cambridge, MA 02139		10. Work Unit No.	
		11. Contract or Grant No. NASW-3950	
		13. Type of Report and Period Covered Interim Report June - August 1985	
12. Sponsoring Agency Name and Address NASA Headquarters Headquarters Contract Division Washington, DC 20546		14. Sponsoring Agency Code HW-2	
15. Supplementary Notes			
16. Abstract The analysis of observed brightness profiles for the C ₂ radical in comets with Haser's model implies that the parent scale length varies as the square of the heliocentric distance. The use of new Haser scale lengths for C ₂ and CN determined under this project appears to explain the <u>apparent</u> drop in the C ₂ -to-CN ratio with increasing heliocentric distance.			
17. Key Words (Selected by Author(s)) comets		18. Distribution Statement	
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of Pages 7	22. Price*

*For sale by the Clearinghouse for Federal Scientific and Technical Information, Springfield, Virginia 22151.

I. Program of Research for the First Quarter

Research activities during this first quarter have involved (1) meeting with Dr. Uwe Fink of the University of Arizona with whom a collaboration has begun in order to analyze observations of the spatial distributions of neutral gases and (2) a preliminary draft of a paper on the observed distributions of the C_2 radical in comets.

1. Model Analysis of New Observations

A collaboration with Dr. Uwe Fink of the University of Arizona had been initiated during the third quarter of the previous project year in which we would analyze the spatial distributions of neutral gases as observed in Dr. Fink's program of CCD long-slit spectrophotometry and narrow-band imaging of comets. During a visit I made to Tucson this past quarter, in addition to my presenting a colloquium on modeling the neutral coma, we previewed raw CCD filtered images and spectra of Comet P/Giacobini-Zinner taken the previous month by Dr. Fink at the 61-inch telescope at Catalina Mountain. Images in the band containing CN emission were clearly distinctive from nearby continuum images, and a likely detection of an [OI] image at 6300 Å was also found.

2. The Distribution and Production of Cometary C_2

A preliminary draft of a paper which deals with the production and observed distribution of C_2 in comets is in preparation. The paper reports new observational data which had been reduced prior to the start of this project. These data are being re-analyzed and combined with other newly-available data in the literature under this project. This work has involved an ongoing collaboration with Dr. A.H. Delsemme of the University of Toledo. The results of this work will also be presented at the Division of Planetary Sciences meeting in Baltimore on October 29, 1985.

The principal results of the paper are (1) new Haser model scale length dependencies on heliocentric distance (see Figures 1 and 2), (2) specification of the parent and C_2 lifetimes, and parent and C_2 ejection velocities determined using the radiation pressure distorted profiles and the Monte Carlo particle-trajectory model, and (3) a resolution of the previously asserted drop of C_2 production in comets relative to other species ($r > 1.2$ AU), which is only an artifact of an incorrect parent scale length law (see Figure 3).




Figure 1. Haser Model Scale Lengths for CN.

The Haser model parent (a) and daughter (b) scale lengths fitted to full observed brightness profiles are plotted as a function of heliocentric distance. In (a) an $r^{1.44}$ law is the best fit to the data and is consistent with the $r^{1.5}$ which would be expected from photodissociation and an $r^{-0.5}$ velocity law. In (b) no observable trend is seen. Data were from Combi (1978), Combi and Delsemme (1980), Cochran (1982), Delsemme and Combi (1983), Johnson et al. (1984).





Figure 2. Haser Model Scale Lengths for C₂.

The Haser model parent (a) and daughter (b) scale lengths fitted to full observed brightness profiles are plotted as a function of heliocentric distance. An $r^{2.0}$ law is the best fit to both sets of scale lengths. The r^2 law for the C₂ parent may mean that the C₂ parent may in fact be a daughter radical. Data are a combination of new data with those from O'Dell and Osterbrock (1962), Delsemme and Miller (1971), Kumar and Southal (1976), Combi (1978), and Cochran (1985).



C_2/CN Ratio in Comet West




Figure 3. The Relative Productions of C_2 and CN in Comet West (1976VI)

The photometry of A'Hearn, Thurber and Millis (1977) is shown as reduced with the Haser model scale length laws assumed by A'Hearn and Cowan (1980) [□] and with the new values found by us [•]. A'Hearn and Cowan used an r^2 law for CN, CN-parent and C_2 , and an r^1 law for the C_2 parent. When we use $r^{1.5}$ for the CN-parent and r^2 for the C_2 parent, the anomalous drop in C_2 production for $r > 1.2$ AU disappears.

II. Program of Research for the Second Quarter

Research activities for the second quarter will concentrate on completion of the C_2 paper with Armand Delsemme and evaluation of the observed C_3 and OH distributions in comets.

REFERENCES

- A'Hearn, M.F. and J.J. Cowan (1980) The Moon and Planets 23, 41.
- A'Hearn, M.F., C.H. Thurber and R.L. Millis (1977) A. J. 82, 518.
- Cochran, A. (1982) Ph.D. thesis, University of Texas.
- Cochran, A. (1985) Ap. J. 289, 388.
- Combi, M.R. (1978) A. J. 83, 1459.
- Combi, M.R., and A.H. Delsemme (1980) Ap. J. 237, 641.
- Delsemme, A.H., and M.R. Combi (1983) Ap. J. 271, 388.
- Delsemme, A.H., and D.C. Miller (1971) Planet. Sp. Sci. 19, 1259.
- Johnson, J.R., U. Fink and S.M. Larson (1984) Icarus 60, 351.
- Kumar, C.K., and R.J. Southal (1966) Proc. IAU Colloq. No. 25, NASA SP-393, p.121.
- O'Dell, C.R., and D.E. Osterbrock (1962) Ap. J. 136, 559.